European Journal of Soil Science, December 2000, **51**, 729–731

European Journal of Soil Science

Volume 51 2000

Author index

Andreu, V. *see* Gimeno-García, E.
Ashman, M.R. *see* Watts, C.W.
Azontonde, A. *see* Barthès, B.

Balesdent, J. *see* Puget, P.
Ball, B.C., Horgan, G.W. & Parker, J.P. Short-range spatial variation of nitrous oxide fluxes in relation to compaction and straw residues, 607
Bardgett, R.D. *see* Cole, L.
Barrow, N.J., Hansen, H.C.B., Hansen, P.E. & Magid, J. A note on the description of the kinetics of phosphate sorption, 531
Barthès, B., Azontonde, A., Boli, B.Z., Prat, C. & Roose, E. Field-scale run-off and erosion in relation to topsoil aggregate stability in three tropical regions (Benin, Cameroon, Mexico), 485
Beaumont, A. *see* Michel, J.-C.
Bengough, A.G. *see* Czarnes, S.
Benoit, P. & Preston, C.M. Transformation and binding of ^{13}C and ^{14}C-labelled atrazine in relation to straw decomposition in soil, 43
Biesheuvel, P.M., van Loon, A.P., Raangs, R., Verweij, H. & Dirksen, C. A prototype osmotic tensiometer with polymeric gel grains, 355
Bird, N.R.A. *see* Watts, C.W.
Bird, N.R.A., Perrier, E. & Rieu, M. The water retention function for a model of soil structure with pore and solid fractal distributions, 55
Blake, L., Goulding, K.W.T., Mott, C.J.B. & Poulton, P.R. Temporal changes in chemical properties of air-dried stored soils and their interpretation for long-term experiments, 345
Blume, H.-P. *see* Wu, Q.
Bœuf-Tremblay, V. *see* Traoré, O.
Boiffin, J. *see* Roger-Estrade, J.
Boizard, H. *see* Roger-Estrade, J.
Boli, B.Z. *see* Barthès, B.
Bourgeon, G. *see* Caner, L.
Brahy, V., Deckers, J. & Delvaux, B. Estimation of soil weathering stage and acid neutralizing capacity in a toposequence Luvisol–Cambisol on loess under deciduous forest in Belgium, 1
Brahy, V., Titeux, H. & Delvaux, B. Incipient podzolization and weathering caused by complexation in a forest Cambisol on loess as revealed by a soil solution study, 475
Brahy, V., Titeux, H., Iserentant, A. & Delvaux, B. Surface podzolization in Cambisols under deciduous forest in the Belgian loess belt, 15
Bruand, A. & Tessier, D. Water retention properties of the clay in soils developed on clayey sediments: significance of parent material and soil history, 679
Brus, D.J. Using regression models in design-based estimation of spatial means of soil properties, 159
Busto, M.D. & Perez-Mateos, M. Characterization of β-D-glucosidase extracted from soil fractions, 193
Buurman, P. *see* Roscoe, R.

Caneill, J. *see* Roger-Estrade, J.
Caner, L., Bourgeon, G., Toutain, F. & Herbillon, A.J. Characteristics of non-allophanic Andisols derived from low-activity clay regoliths in the Nilgiri Hills (Southern India), 553
Carstensen, A. *see* Young, S.D.
Chadœuf, J. *see* Parry, S.
Chenu, C. *see* Parry, S.; Puget, P.
Cheshire, M.V., Dumat, C., Fraser, A.R., Hillier, S. & Staunton, S. The interaction between soil organic matter and soil clay minerals by selective removal and controlled addition of organic matter, 497
Choi, S.I. *see* Kim, D.J.
Christl, I., Knicker, H., Kögel-Knabner, I. & Kretzschmar, R. Chemical heterogeneity of humic substances: characterization of size fractions obtained by hollow-fibre ultrafiltration, 617
Cole, L., Bardgett, R.D. & Ineson, P. Enchytraeid worms (Oligochaeta) enhance mineralization of carbon in organic upland soils, 185
Coleman, K. *see* Romanyà, J.
Conen, F. & Smith, K.A. An explanation of linear increases in gas concentration under closed chambers used to measure gas exchange between soil and the atmosphere, 111
Conrad, R. *see* Yao, H.
Cortina, J. *see* Romanyà, J.
Crout, N. *see* Young, S.D.
Czarnes, S., Hallett, P.D., Bengough, A.G. & Young, I.M. Root- and microbial-derived mucilages affect soil structure and water transport, 435

Da Silva, L.F. *see* Schroth, G.
Darrah, P.R. & Staunton, S. A mathematical model of root uptake of cations incorporating root turnover, distribution within the plant, and recycling of absorbed species, 643
De Neve, S., Van de Steene, J., Hartmann, R. & Hofman, G. Using time domain reflectometry for monitoring mineralization of nitrogen from soil organic matter, 295
De Smedt, F. *see* Van Praag, H.J.
Deckers, J. *see* Brahy, V.
Delvaux, B. *see* Brahy, V.
Descamps, M. *see* Grelle, C.
Dirksen, C. *see* Biesheuvel, P.M.
Dumat, C. *see* Cheshire, M.V.

Elliott, E.T. *see* Six, J.

Fabre, M.-C. *see* Grelle, C.
Falloon, P. *see* Romanyà, J.
Feyen, J. *see* Kim, D.J.
Flühler, H. *see* Forrer, I.
Fölschow, M. *see* Wu, Q.
Forrer, I., Papritz, A., Kasteel, R., Flühler, H. & Luca, D. Quantifying dye tracers in soil profiles by image processing, 313
Fraser, A.R. *see* Cheshire, M.V.

© 2000 Blackwell Science Ltd

Fritze, H., Pietikäinen, J. & Pennanen, T. Distribution of microbial biomass and phospholipid fatty acids in Podzol profiles under coniferous forest, 565

Germon, J.C. *see* Hénault, C.
Gerzabek, M.H. *see* Spaccini, R.
Gillman, G.P. *see* Noble, A.D.
Gimeno-García, E., Andreu, V. & Rubio, J.L. Changes in organic matter, nitrogen, phosphorus and cations in soil as a result of fire and water erosion in a Mediterranean landscape, 201
Goulding, K.W.T. *see* Blake, L.
Grelle, C., Fabre, M.-C., Leprêtre, A. & Descamps, M. Myriapod and isopod communities in soils contaminated by heavy metals in northern France, 425
Groleau-Renaud, V. *see* Traoré, O.

Haberhauer, G. *see* Spaccini, R.
Hallett, P.D. *see* Czarnes, S.
Hansen, H.C.B. *see* Barrow, N.J.
Hansen, P.E. *see* Barrow, N.J.
Hartmann, R. *see* De Neve, S.
Hénault, C. & Germon, J.C. NEMIS, a predictive model of denitrification on the field scale, 257
Herbillon, A.J. *see* Caner, L.
Hillier, S. *see* Cheshire, M.V.
Hiraiwa, Y. & Kasubuchi, T. Temperature dependence of thermal conductivity of soil over a wide range of temperature (5–75°C), 211
Hirsch, C. *see* Prietzel, J.
Hofman, G. *see* De Neve, S.
Hongve, D., van Hees, P.A.W. & Lundström, U.S. Dissolved components in precipitation water percolated through forest litter, 667
Horgan, G.W. *see* Ball, B.C.

Ineson, P. *see* Cole, L.
Iserentant, A. *see* Brahy, V.

Jaakkola, A. *see* Simojoki, A.
Janssen, B.H. *see* Yang, H.S.
Jarvis, S.C. *see* Peat, D.M.W.
Joel, A. & Messing, I. Application of two methods to determine hydraulic conductivity with disc permeameters on sloping land, 93

Kaiser, K. & Zech, W. Sorption of dissolved organic nitrogen by acid subsoil horizons and individual mineral phases, 403
Kasteel, R. *see* Forrer, I.
Kasteel, R., Vogel, H.-J. & Roth, K. From local hydraulic properties to effective transport in soil, 81
Kasubuchi, T. *see* Hiraiwa, Y.
Kim, D.J., Choi, S.I., Ryszard, O., Feyen, J. & Kim, H.S. Determination of moisture content in a deformable soil using time-domain reflectometry (TDR), 119
Kim, H.S. *see* Kim, D.J.
Kimpe, K. *see* Six, J.
Knicker, H. *see* Christl, I.
Kögel-Knabner, I. *see* Christl, I.
Kretzschmar, R. *see* Christl, I.

Lark, R.M. A comparison of some robust estimators of the variogram for use in soil survey, 137
Lark, R.M. Estimating variograms of soil properties by the method-of-moments and maximum likelihood, 717
Lehmann, J. *see* Solomon, D.
Lensi, R. *see* Parry, S.
Leprêtre, A. *see* Grelle, C.
Loch, J.P.G. *see* Van den Berg, G.A.
Luca, D. *see* Forrer, I.
Lundström, U.S. *see* Hongve, D.

Magid, J. *see* Barrow, N.J.
Mahieu, N., Olk, D.C. & Randall, E.W. Accumulation of heterocyclic nitrogen in humified organic matter: a ^{15}N-NMR study of lowland rice soils, 379
Mahieu, N., Olk, D.C. & Randall, E.W. Analysis of phosphorus in two humic acid fractions of intensively cropped lowland rice soils by ^{31}P-NMR, 391
Manichon, H. *see* Roger-Estrade, J.
Matthews, G.P. *see* Peat, D.M.W.
Merckx, R. *see* Six, J.
Messing, I. *see* Joel, A.
Michel, J.-C., Beaumont, A. & Tessier, D. A laboratory method for measuring the isotropic character of soil swelling, 689
Morel, C. *see* Schneider, A.
Mott, C.J.B. *see* Blake, L.

Noble, A.D., Gillman, G.P. & Ruaysoongnern, S. A cation exchange index for assessing degradation of acid soil by further acidification under permanent agriculture in the tropics, 233

Olk, D.C. *see* Mahieu, N.

Papritz, A. *see* Forrer, I.
Parfitt, R.L. *see* Yuan, G.
Parker, J.P. *see* Ball, B.C.
Parry, S., Renault, P., Chadœuf, J., Chenu, C. & Lensi, R. Particulate organic matter as a source of variation in denitrification in clods of soil, 271
Paustian, K. *see* Six, J.
Peat, D.M.W., Matthews, G.P., Worsfold, P.J. & Jarvis, S.C. Simulation of water retention and hydraulic conductivity in soil using a three-dimensional network, 65
Pennanen, T. *see* Fritze, H.
Percival, H.J. *see* Yuan, G.
Perez-Mateos, M. *see* Busto, M.D.
Perrier, E. *see* Bird, N.R.A.
Piccolo, A. *see* Spaccini, R.
Pietikäinen, J. *see* Fritze, H.
Plantureux, S. *see* Traoré, O.
Poulton, P.R. *see* Blake, L.
Prat, C. *see* Barthès, B.
Preston, C.M. *see* Benoit, P.
Prietzel, J. & Hirsch, C. Ammonium fluoride extraction for determining inorganic sulphur in acid forest soils, 323
Puget, P., Chenu, C. & Balesdent, J. Dynamics of soil organic matter associated with particle-size fractions of water-stable aggregates, 595

Raangs, R. *see* Biesheuvel, P.M.
Randall, E.W. *see* Mahieu, N.
Rayment, M.B. Closed chamber systems underestimate soil CO_2 efflux, 107
Renault, P. *see* Parry, S.
Resende, L. *see* Young, S.D.
Rexilius, L. *see* Wu, Q.
Richard, G. *see* Roger-Estrade, J.
Rieu, M. *see* Bird, N.R.A.
Rimmer, D.L. *see* Uygur, V.
Roger-Estrade, J., Richard, G., Boizard, H., Boiffin, J., Caneill, J. & Manichon, H. Modelling structural changes in tilled topsoil over time as a function of cropping systems, 455
Romanyà, J., Cortina, J., Falloon, P., Coleman, K. & Smith, P. Modelling changes in soil organic matter after planting fast-growing *Pinus radiata* on Mediterranean agricultural soils, 627
Roose, E. *see* Barthès, B.
Roscoe, R., Buurman, P. & Velthorst, E.J. Disruption of soil aggregates by varied amounts of ultrasonic energy in fractionation of organic matter of a clay Latosol: carbon, nitrogen and δ^{13}C distribution in particle-size fractions, 445
Roth, K. *see* Kasteel, R.
Ruaysoongnern, S. *see* Noble, A.D.
Rubio, J.L. *see* Gimeno-García, E.
Ryszard, O. *see* Kim, D.J.

Schleuss, U. *see* Wu, Q.

Schneider, A. & Morel, C. Relationship between the isotopically exchangeable and resin-extractable phosphate of deficient to heavily fertilized soil, 709
Schroth, G., Seixas, R., da Silva, L.F., Teixeira, W.G. & Zech, W. Nutrient concentrations and acidity in ferralitic soil under perennial cropping, fallow and primary forest in central Amazonia, 219
Seixas, R. *see* Schroth, G.
Simojoki, A. & Jaakkola, A. Effect of nitrogen fertilization, cropping and irrigation on soil air composition and nitrous oxide emission in a loamy clay, 413
Simonsson, M. Interactions of aluminium and fulvic acid in moderately acid solutions: stoichiometry of the H^+/Al^{3+} exchange, 655
Six, J., Merckx, R., Kimpe, K., Paustian, K. & Elliott, E.T. A re-evaluation of the enriched labile soil organic matter fraction, 283
Smith, K.A. *see* Conen, F.
Smith, P. *see* Romanyà, J.
Solomon, D. & Lehmann, J. Loss of phosphorus from soil in semi-arid northern Tanzania as a result of cropping: evidence from sequential extraction and ^{31}P-NMR spectroscopy, 699
Spaccini, R., Piccolo, A., Haberhauer, G. & Gerzabek, M.H. Transformation of organic matter from maize residues into labile and humic fractions of three European soils as revealed by ^{13}C distribution and CPMAS-NMR spectra, 583
Staunton, S. *see* Cheshire, M.V.; Darrah, P.R.

Teixeira, W.G. *see* Schroth, G.
Tessier, D. *see* Bruand, A.; Michel, J.-C.
Theng, B.K.G. *see* Yuan, G.
Titeux, H. *see* Brahy, V.
Toutain, F. *see* Caner, L.
Traoré, O., Groleau-Renaud, V., Plantureux, S., Tubeileh, A. & Bœuf-Tremblay, V. Effect of root mucilage and modelled root exudates on soil structure, 575
Tubeileh, A. *see* Traoré, O.
Tye, A. *see* Young, S.D.

Uygur, V. & Rimmer, D.L. Reactions of zinc with iron-oxide coated calcite surfaces at alkaline pH, 511

Van de Steene, J. *see* De Neve, S.
Van den Berg, G.A. & Loch, J.P.G. Decalcification of soils subject to periodic waterlogging, 27
Van Hees, P.A.W. *see* Hongve, D.
Van Loon, A.P. *see* Biesheuvel, P.M.
Van Praag, H.J., De Smedt, F. & vu Thanh, T. Simulation of calcium leaching and desorption in an acid forest soil, 245
Velthorst, E.J. *see* Roscoe, R.
Verweij, H. *see* Biesheuvel, P.M.
Vogel, H.J. A numerical experiment on pore size, pore connectivity, water retention, permeability, and solute transport using network models, 99
Vogel, H.-J. *see* Kasteel, R.
Vu Thanh, T. *see* Van Praag, H.J.

Watts, C.W., Whalley, W.R., Bird, N.R.A. & Ashman, M.R. The effect of iron concentration, hindered settling, saturating cation and aggregate density of clays on the size distribution determined by gravitation X-ray sedimentometry, 305
Whalley, W.R. *see* Watts, C.W.
Worsfold, P.J. *see* Peat, D.M.W.
Wu, Q., Blume, H.-P., Rexilius, L., Fölschow, M. & Schleuss, U. Sorption of atrazine, 2,4-D, nitrobenzene and pentachlorophenol by urban and industrial wastes, 335

Yang, H.S. & Janssen, B.H. A mono-component model of carbon mineralization with a dynamic rate constant, 517
Yao, H. & Conrad, R. Electron balance during steady-state production of CH_4 and CO_2 in anoxic rice soil, 369
Young, I.M. *see* Czarnes, S.
Young, S.D., Tye, A., Carstensen, A., Resende, L. & Crout, N. Methods for determining labile cadmium and zinc in soil, 129
Yuan, G., Theng, B.K.G., Parfitt, R.L. & Percival, H.J. Interactions of allophane with humic acid and cations, 35

Zech, W. *see* Kaiser, K.; Schroth, G.

European Journal of Soil Science, December 2000, **51**, 732–736

European Journal of Soil Science

Volume 51 2000

Subject index

accumulation of heterocyclic nitrogen in humified organic matter: a ^{15}N-NMR study of lowland rice soils, 379
acid forest soil
  ammonium fluoride extraction for determining inorganic sulphur in, 323
  simulation of calcium leaching and desorption in an, 245
acid neutralizing capacity in a toposequence Luvisol–Cambisol on loess under deciduous forest in Belgium, estimation of, 1
acid subsoil horizons, sorption of dissolved organic nitrogen by, 403
acidification under permanent agriculture in the tropics, cation exchange index for assessing degradation of acid soil by further, 233
acidity in ferralitic soil under perennial cropping, fallow and primary forest in central Amazonia, 219
aggregate density of clays, effect of, on the size distribution determined by gravitation X-ray sedimentometry, 305
agriculture, permanent, in the tropics, cation exchange index for assessing degradation of acid soil by further acidification under, 233
air-dried stored soils, temporal changes in chemical properties of, and their interpretation for long-term experiments, 345
allophane, interactions with humic acid and cations, 35
aluminium and fulvic acid in moderately acid solutions, interactions of: stoichiometry of the H^+/Al^{3+} exchange, 655
Amazonia, central, nutrient concentrations and acidity in ferralitic soil under perennial cropping, fallow and primary forest in, 219
ammonium fluoride extraction for determining inorganic sulphur in acid forest soils, 323
analysis of phosphorus in two humic acid fractions of intensively cropped lowland rice soils by ^{31}P-NMR, 391
Andisols, non-allophanic, derived from low-activity clay regoliths in the Nilgiri Hills (Southern India), characteristics of, 553
anoxic rice soil, electron balance during steady-state production of CH_4 and CO_2 in, 369
application of two methods to determine hydraulic conductivity with disc permeameters on sloping land, 93
atrazine
  sorption of, by urban and industrial wastes, 335
  transformation and binding of ^{13}C and ^{14}C-labelled, in relation to straw decomposition in soil, 43

β-D-glucosidase extracted from soil fractions, characterization of, 193
Belgian loess belt, surface podzolization in Cambisols under deciduous forest in the, 15
Belgium, estimation of soil weathering stage and acid neutralizing capacity in a toposequence Luvisol–Cambisol on loess under deciduous forest in, 1

^{13}C and ^{14}C-labelled atrazine, transformation and binding of, in relation to straw decomposition in soil, 43
^{13}C distribution, transformation of organic matter from maize residues into labile and humic fractions of three European soils as revealed by, 583
cadmium in soil, methods for determining labile, 129
calcite surfaces, iron-oxide coated, reactions of zinc with, at alkaline pH, 511
calcium leaching and desorption in an acid forest soil, simulation of, 245
Cambisols under deciduous forest in the Belgian loess belt, surface podzolization in, 15
carbon
  distribution in particle-size fractions: disruption of soil aggregates by varied amounts of ultrasonic energy in fractionation of organic matter of a clay Latosol, 445
  in organic upland soils, enchytraeid worms (Oligochaeta) enhance mineralization of, 185
  mineralization, mono-component model of, with a dynamic rate constant, 517
cation exchange index for assessing degradation of acid soil by further acidification under permanent agriculture in the tropics, 233
cations
  allophane interactions with, 35
  changes in, as a result of fire and water erosion in a Mediterranean landscape, 201
  mathematical model of root uptake of, incorporating root turnover, distribution within the plant, and recycling of absorbed species, 643
central Amazonia, nutrient concentrations and acidity in ferralitic soil under perennial cropping, fallow and primary forest in, 219
CH_4, electron balance during steady-state production of, in anoxic rice soil, 369
changes in organic matter, nitrogen, phosphorus and cations in soil as a result of fire and water erosion in a Mediterranean landscape, 201
characterization of β-D-glucosidase extracted from soil fractions, 193
chemical heterogeneity of humic substances: characterization of size fractions obtained by hollow-fibre ultrafiltration, 617
chemical properties of air-dried stored soils and their interpretation for long-term experiments, temporal changes in, 345
clay
  in soils developed on clayey sediments, water retention properties of: significance of parent material and soil history, 679
  Latosol, disruption of soil aggregates by varied amounts of ultrasonic energy in fractionation of organic matter of a: carbon, nitrogen and δ^{13}C distribution in particle-size fractions, 445
  regoliths, low-activity, in the Nilgiri Hills (Southern India), characteristics of non-allophanic Andisols derived from, 553
closed chambers
  to measure gas exchange between soil and the atmosphere, explanation of linear increases in gas concentration under, 111
  underestimate soil CO_2 efflux, 107
CO_2
  efflux, from soil, closed chamber systems underestimate, 107
  electron balance during steady-state production of, in anoxic rice soil, 369
compaction, short-range spatial variation of nitrous oxide fluxes in relation to, 607

© 2000 Blackwell Science Ltd

comparison of some robust estimators of the variogram for use in soil survey, 137
complexation, incipient podzolization and weathering caused by, in a forest Cambisol on loess as revealed by a soil solution study, 475
coniferous forest, distribution of microbial biomass and phospholipid fatty acids in Podzol profiles under, 565
CPMAS-NMR spectra, transformation of organic matter from maize residues into labile and humic fractions of three European soils as revealed by, 583
cropping
effect of, on soil air composition and nitrous oxide emission in a loamy clay, 413
loss of phosphorus from soil in semi-arid northern Tanzania as a result of: evidence from sequential extraction and ^{31}P-NMR spectroscopy, 699
systems, modelling structural changes in tilled topsoil over time as a function of, 455

2,4-D, sorption of, by urban and industrial wastes, 335
decalcification of soils subject to periodic waterlogging, 27
deciduous forest
estimation of soil weathering stage and acid neutralizing capacity in a toposequence Luvisol–Cambisol on loess, under, 1
surface podzolization in Cambisols under, 15
degradation of acid soil by further acidification under permanent agriculture in the tropics, cation exchange index for assessing, 233
denitrification
in clods of soil, particulate organic matter as a source of variation in, 271
on the field scale, NEMIS, a predictive model of, 257
description of the kinetics of phosphate sorption, 531
design-based estimation of spatial means of soil properties, regression models in, 159
determination of moisture content in a deformable soil using time-domain reflectometry (TDR), 119
disc permeameters, application of two methods to determine hydraulic conductivity with, on sloping land, 93
disruption of soil aggregates by varied amounts of ultrasonic energy in fractionation of organic matter of a clay Latosol: carbon, nitrogen and δ^{13}C distribution in particle-size fractions, 445
dissolved components in precipitation water percolated through forest litter, 667
dissolved organic nitrogen, sorption of, by acid subsoil horizons and individual mineral phases, 403
distribution
of microbial biomass and phospholipid fatty acids in Podzol profiles under coniferous forest, 565
within the plant, mathematical model of root uptake of cations incorporating, 643
dye tracers in soil profiles by image processing, quantifying, 313
dynamic rate constant, mono-component model of carbon mineralization with a, 517

effect
of iron concentration, hindered settling, saturating cation and aggregate density of clays on the size distribution determined by gravitation X-ray sedimentometry, 305
of nitrogen fertilization, cropping and irrigation on soil air composition and nitrous oxide emission in a loamy clay, 413
effective transport in soil, to, from local hydraulic properties, 81
electron balance during steady-state production of CH_4 and CO_2 in anoxic rice soil, 369
enchytraeid worms (Oligochaeta) enhance mineralization of carbon in organic upland soils, 185
enriched labile soil organic matter fraction, re-evaluation of the, 283
erosion
by fire and water, changes in organic matter, nitrogen, phosphorus and cations in soil as a result of, in a Mediterranean landscape, 201
field-scale, in relation to topsoil aggregate stability in three tropical regions (Benin, Cameroon, Mexico), 485
estimation of soil weathering stage and acid neutralizing capacity in a toposequence Luvisol–Cambisol on loess under deciduous forest in Belgium, 1
European soils, transformation of organic matter from maize residues into labile and humic fractions of three, as revealed by ^{13}C distribution and CPMAS-NMR spectra, 583
explanation of linear increases in gas concentration under closed chambers used to measure gas exchange between soil and the atmosphere, 111

ferralitic soil under perennial cropping, fallow and primary forest in central Amazonia, nutrient concentrations and acidity in, 219
field-scale run-off and erosion in relation to topsoil aggregate stability in three tropical regions (Benin, Cameroon, Mexico), 485
fire and water erosion, changes in organic matter, nitrogen, phosphorus and cations in soil as a result of, in a Mediterranean landscape, 201
forest
Cambisol on loess, incipient podzolization and weathering caused by complexation in a, as revealed by a soil solution study, 475
litter, dissolved components in precipitation water percolated through, 667
fractal distributions, pore and solid, water retention function for a model of soil structure with, 55
fractionation of organic matter of a clay Latosol, disruption of soil aggregates by varied amounts of ultrasonic energy in: carbon, nitrogen and δ^{13}C distribution in particle-size fractions, 445
France, northern, myriapod communities in soils contaminated by heavy metals in, 425

gas
concentration, explanation of linear increases in, under closed chambers used to measure gas exchange between soil and the atmosphere, 111
exchange between soil and the atmosphere, explanation of linear increases in gas concentration under closed chambers used to measure, 111
gravitation X-ray sedimentometry, effect of iron concentration, hindered settling, saturating cation and aggregate density of clays on the size distribution determined by, 305

H^+/Al^{3+} exchange, stoichiometry of, in moderately acid solutions, 655
heavy metals in northern France, myriapod and isopod communities in soils contaminated by, 425
heterocyclic nitrogen, accumulation of, in humified organic matter: a ^{15}N-NMR study of lowland rice soils, 379
hindered settling, effect of, on the size distribution determined by gravitation X-ray sedimentometry, 305
hollow-fibre ultrafiltration, characterization of size fractions of humic substances obtained by, 617
humic acid
allophane interactions with, 35
fractions of intensively cropped lowland rice soils, analysis of phosphorus in two, by ^{31}P-NMR, 391
humic substances, chemical heterogeneity of: characterization of size fractions obtained by hollow-fibre ultrafiltration, 617
humified organic matter, accumulation of heterocyclic nitrogen in: a ^{15}N-NMR study of lowland rice soils, 379
hydraulic conductivity
application of two methods to determine, with disc permeameters on sloping land, 93
simulation of, using a three-dimensional network, 65

image processing, quantifying dye tracers in soil profiles by, 313
India, Southern, characteristics of non-allophanic Andisols derived from low-activity clay regoliths in the Nilgiri Hills in, 553
individual mineral phases, sorption of dissolved organic nitrogen by, 403
interaction
between soil organic matter and soil clay minerals by selective removal and controlled addition of organic matter, 497
of allophane with humic acid and cations, 35
iron concentration, effect of, on the size distribution determined by gravitation X-ray sedimentometry, 305
iron-oxide coated calcite surfaces at alkaline pH, reactions of zinc with, 511
irrigation, effect of, on soil air composition and nitrous oxide emission in a loamy clay, 413

isopod communities in soils contaminated by heavy metals in northern France, 425
isotopically exchangeable phosphate, relationship between, and resin-extractable phosphate of deficient to heavily fertilized soil, 709
isotropic character of soil swelling, a laboratory method for measuring the, 689

kinetics of phosphate sorption, description of the, 531

labile
and humic fractions of three European soils, transformation of organic matter from maize residues into, as revealed by ^{13}C distribution and CPMAS-NMR spectra, 583
cadmium in soil, methods for determining, 129
soil organic matter fraction, enriched, re-evaluation of the, 283
zinc in soil, methods for determining, 129
laboratory method for measuring the isotropic character of soil swelling, 689
Latosol, clay, disruption of soil aggregates by varied amounts of ultrasonic energy in fractionation of organic matter of a: carbon, nitrogen and $\delta^{13}C$ distribution in particle-size fractions, 445
loamy clay, effect of nitrogen fertilization, cropping and irrigation on soil air composition and nitrous oxide emission in a, 413
local hydraulic properties, from, to effective transport in soil, 81
loess
estimation of soil weathering stage and acid neutralizing capacity in a toposequence Luvisol–Cambisol on, under deciduous forest in Belgium, I
incipient podzolization and weathering caused by complexation in a forest Cambisol on, as revealed by a soil solution study, 475
long-term experiments, temporal changes in chemical properties of air-dried stored soils and their interpretation for, 345
lowland rice soils
accumulation of heterocyclic nitrogen in humified organic matter: a ^{15}N-NMR study of, 379
analysis of phosphorus in two humic acid fractions of intensively cropped, by ^{31}P-NMR, 391

mathematical model of root uptake of cations incorporating root turnover, distribution within the plant, and recycling of absorbed species, 643
maximum likelihood, estimating variograms of soil properties by, 717
Mediterranean
agricultural soils, modelling changes in soil organic matter after planting fast-growing *Pinus radiata* on, 627
landscape, changes in organic matter, nitrogen, phosphorus and cations in soil as a result of fire and water erosion in a, 201
method-of-moments, estimating variograms of soil properties by, 717
methods for determining labile cadmium and zinc in soil, 129
microbial biomass, distribution of, in Podzol profiles under coniferous forest, 565
mineralization
of carbon in organic upland soils, enhancement of, by enchytraeid worms (Oligochaeta), 185
of nitrogen from soil organic matter, monitoring using time domain reflectometry, 295
model
mono-component, of carbon mineralization with a dynamic rate constant, 517
of soil structure, water retention function for, with pore and solid fractal distributions, 55
modelled root exudates, effect of, on soil structure, 575
modelling structural changes in tilled topsoil over time as a function of cropping systems, 455
models, regression, in design-based estimation of spatial means of soil properties, 159
moisture content in a deformable soil, determination of, using time-domain reflectometry (TDR), 119
mono-component model of carbon mineralization with a dynamic rate constant, 517
mucilages, root- and microbial-derived, affect soil structure and water transport, 435

myriapod communities in soils contaminated by heavy metals in northern France, 425

^{15}N-NMR study of lowland rice soils: accumulation of heterocyclic nitrogen in humified organic matter, 379
NEMIS, a predictive model of denitrification on the field scale, 257
network models, numerical experiment on pore size, pore connectivity, water retention, permeability, and solute transport using, 99
nitrobenzene, sorption of, by urban and industrial wastes, 335
nitrogen
changes in, as a result of fire and water erosion in a Mediterranean landscape, 201
dissolved organic, sorption of, by acid subsoil horizons and individual mineral phases, 403
fertilization, effect of, on soil air composition and nitrous oxide emission in a loamy clay, 413
heterocyclic, accumulation of, in humified organic matter: a ^{15}N-NMR study of lowland rice soils, 379
monitoring mineralization of, from soil organic matter using time domain reflectometry, 295
nitrous oxide
emission in a loamy clay, effect of nitrogen fertilization, cropping and irrigation on, 413
fluxes, short-range spatial variation of, in relation to compaction and straw residues, 607
non-allophanic Andisols derived from low-activity clay regoliths in the Nilgiri Hills (Southern India), characteristics of, 553
northern France, myriapod communities in soils contaminated by heavy metals in, 425
numerical experiment on pore size, pore connectivity, water retention, permeability, and solute transport using network models, 99
nutrient concentrations and acidity in ferralitic soil under perennial cropping, fallow and primary forest in central Amazonia, 219

organic matter
changes in, as a result of fire and water erosion in a Mediterranean landscape, 201
from maize residues, transformation of, into labile and humic fractions of three European soils as revealed by ^{13}C distribution and CPMAS-NMR spectra, 583
interaction between soil organic matter and soil clay minerals by selective removal and controlled addition of, 497
of a clay Latosol, disruption of soil aggregates by varied amounts of ultrasonic energy in fractionation of: carbon, nitrogen and $\delta^{13}C$ distribution in particle-size fractions, 445
particulate, as a source of variation in denitrification in clods of soil, 271
osmotic tensiometer, prototype, with polymeric gel grains, 355

^{31}P-NMR
evidence from, for loss of phosphorus from soil in semi-arid northern Tanzania as a result of cropping, 699
analysis of phosphorus in two humic acid fractions of intensively cropped lowland rice soils by, 391
parent material, significance of, on water retention properties of the clay in soils developed on clayey sediments, 679
particle-size fractions
carbon, nitrogen and $\delta^{13}C$ distribution in: disruption of soil aggregates by varied amounts of ultrasonic energy in fractionation of organic matter of a clay Latosol, 445
of water-stable aggregates, dynamics of soil organic matter associated with, 595
particulate organic matter as a source of variation in denitrification in clods of soil, 271
pentachlorophenol, sorption of, by urban and industrial wastes, 335
perennial cropping, fallow and primary forest in central Amazonia, nutrient concentrations and acidity in ferralitic soil under, 219
periodic waterlogging, decalcification of soils subject to, 27
permeability, numerical experiment on using network models, 99
phosphate
of deficient to heavily fertilized soil, relationship between isotopically exchangeable and resin-extractable forms, 709
sorption, description of the kinetics of, 531

phospholipid fatty acids, distribution of, in Podzol profiles under coniferous forest, 565
phosphorus
analysis of, in two humic acid fractions of intensively cropped lowland rice soils by ^{31}P-NMR, 391
changes in, as a result of fire and water erosion in a Mediterranean landscape, 201
loss of, from soil in semi-arid northern Tanzania as a result of cropping: evidence from sequential extraction and ^{31}P-NMR spectroscopy, 699
Pinus radiata on Mediterranean agricultural soils, modelling changes in soil organic matter after planting fast-growing, 627
Podzol profiles under coniferous forest, distribution of microbial biomass and phospholipid fatty acids in, 565
podzolization
incipient, caused by complexation in a forest Cambisol on loess as revealed by a soil solution study, 475
surface, in Cambisols under deciduous forest in the Belgian loess belt, 15
polymeric gel grains, prototype osmotic tensiometer with, 355
pore
connectivity, numerical experiment on using network models, 99
fractal distributions, water retention function for a model of soil structure with, 55
size, numerical experiment on using network models, 99
precipitation water percolated through forest litter, dissolved components in, 667
prototype osmotic tensiometer with polymeric gel grains, 355

quantifying dye tracers in soil profiles by image processing, 313

reactions of zinc with iron-oxide coated calcite surfaces at alkaline pH, 511
recycling of absorbed species, mathematical model of root uptake of cations incorporating, 643
re-evaluation of the enriched labile soil organic matter fraction, 283
regression models in design-based estimation of spatial means of soil properties, 159
resin-extractable phosphate, relationship between, and isotopically exchangeable phosphate of deficient to heavily fertilized soil, 709
rice soil
anoxic, electron balance during steady-state production of CH_4 and CO_2 in, 369
rice soils, lowland
accumulation of heterocyclic nitrogen in humified organic matter: a ^{15}N-NMR study of, 379
analysis of phosphorus in two humic acid fractions of intensively cropped, by ^{31}P-NMR, 391
robust estimators of the variogram for use in soil survey, comparison of some, 137
root- and microbial-derived mucilages affect soil structure and water transport, 435
root
mucilage, effect of, on soil structure, 575
turnover, mathematical model of root uptake of cations incorporating, 643
uptake of cations, mathematical model of, incorporating root turnover, distribution within the plant, and recycling of absorbed species, 643
run-off, field-scale, in relation to topsoil aggregate stability in three tropical regions (Benin, Cameroon, Mexico), 485

saturating cation, effect of, on the size distribution determined by gravitation X-ray sedimentometry, 305
sequential extraction, evidence from, for loss of phosphorus from soil in semi-arid northern Tanzania as a result of cropping, 699
simulation
of calcium leaching and desorption in an acid forest soil, 245
of water retention and hydraulic conductivity in soil using a three-dimensional network, 65
size distribution determined by gravitation X-ray sedimentometry, effect of iron concentration, hindered settling, saturating cation and aggregate density of clays on the, 305
size fractions of humic substances obtained by hollow-fibre ultrafiltration, chemical heterogeneity and characterization of, 617
soil aggregates, disruption of, by varied amounts of ultrasonic energy in fractionation of organic matter of a clay Latosol: carbon, nitrogen and δ^{13}C distribution in particle-size fractions, 445
soil air composition in a loamy clay, effect of nitrogen fertilization, cropping and irrigation on, 413
soil CO_2 efflux, closed chamber systems underestimate, 107
soil fractions, characterization of β-D-glucosidase extracted from, 193
soil history, significance of, on water retention properties of the clay in soils developed on clayey sediments, 679
soil organic matter
and soil clay minerals, interaction between, by selective removal and controlled addition of organic matter, 497
associated with particle-size fractions of water-stable aggregates, dynamics of, 595
fraction, re-evaluation of the enriched labile, 283
modelling changes in, after planting fast-growing *Pinus radiata* on Mediterranean agricultural soils, 627
time domain reflectometry for monitoring mineralization of nitrogen from, 295
soil properties
regression models in design-based estimation of spatial means of, 159
variograms of, estimating by the method-of-moments and maximum likelihood, 717
soil structure
effect of root mucilage and modelled root exudates on, 575
root- and microbial-derived mucilages affect, 435
soil survey, comparison of some robust estimators of the variogram for use in, 137
soil swelling, the isotropic character of, a laboratory method for measuring, 689
soil weathering stage in a toposequence Luvisol–Cambisol on loess under deciduous forest in Belgium, estimation of, 1
solid fractal distributions, water retention function for a model of soil structure with, 55
solute transport, numerical experiment on using network models, 99
sorption
of atrazine, 2,4-D, nitrobenzene and pentachlorophenol by urban and industrial wastes, 335
of dissolved organic nitrogen by acid subsoil horizons and individual mineral phases, 403
steady-state production of CH_4 and CO_2 in anoxic rice soil, electron balance during, 369
straw
decomposition in soil, transformation and binding of ^{13}C and ^{14}C-labelled atrazine in relation to, 43
residues, short-range spatial variation of nitrous oxide fluxes in relation to, 607
structural changes in tilled topsoil over time as a function of cropping systems, modelling, 455
sulphur, inorganic, in acid forest soils, ammonium fluoride extraction for determining inorganic, 323
surface podzolization in Cambisols under deciduous forest in the Belgian loess belt, 15

Tanzania, semi-arid northern, loss of phosphorus from soil in, as a result of cropping: evidence from sequential extraction and ^{31}P-NMR spectroscopy, 699
temperature dependence of thermal conductivity of soil over a wide range of temperature (5–75°C), 211
temporal changes in chemical properties of air-dried stored soils and their interpretation for long-term experiments, 345
thermal conductivity of soil over a wide range of temperature (5–75°C), temperature dependence of, 211
three-dimensional network, simulation of water retention and hydraulic conductivity in soil using a, 65
tilled topsoil, modelling structural changes in, over time as a function of cropping systems, 455
time domain reflectometry
determination of moisture content in a deformable soil using, 119
for monitoring mineralization of nitrogen from soil organic matter, 295

toposequence Luvisol–Cambisol on loess under deciduous forest in Belgium, estimation of soil weathering stage and acid neutralizing capacity in, 1
topsoil aggregate stability, field-scale run-off and erosion in relation to, in three tropical regions (Benin, Cameroon, Mexico), 485
transformation and binding of ^{13}C and ^{14}C-labelled atrazine in relation to straw decomposition in soil, 43
tropical regions, field-scale run-off and erosion in relation to topsoil aggregate stability in three (Benin, Cameroon, Mexico), 485
tropics, cation exchange index for assessing degradation of acid soil by further acidification under permanent agriculture in the, 233

ultrasonic energy, disruption of soil aggregates by varied amounts of, in fractionation of organic matter of a clay Latosol: carbon, nitrogen and $\delta^{13}C$ distribution in particle-size fractions, 445
upland soils, organic, enchytraeid worms (Oligochaeta) enhance mineralization of carbon in, 185
urban and industrial wastes, sorption of atrazine, 2,4-D, nitrobenzene and pentachlorophenol by, 335

variogram for use in soil survey, comparison of some robust estimators of the, 137
variograms of soil properties, estimating by the method-of-moments and maximum likelihood, 717

water retention
  function for a model of soil structure with pore and solid fractal distributions, 55
  in soil, simulation of, using a three-dimensional network, 65
  numerical experiment on using network models, 99
  properties of the clay in soils developed on clayey sediments: significance of parent material and soil history, 679
water transport, root- and microbial-derived mucilages affect, 435
waterlogging, decalcification of soils subject to periodic, 27
water-stable aggregates, dynamics of soil organic matter associated with particle-size fractions of, 595
weathering, caused by complexation in a forest Cambisol on loess as revealed by a soil solution study, 475
worms, enchytraeid, (Oligochaeta) enhance mineralization of carbon in organic upland soils, 185

zinc
  in soil, methods for determining labile, 129
  reactions of, with iron-oxide coated calcite surfaces at alkaline pH, 511